

ENTOCORT®

Budesonide

Consumer Medicine Information

Name of consumer

Address of consumer

What is in this leaflet

This leaflet answers some common questions about Entocort.

It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking Entocort against the benefits they expect it will have for you.

If you have any concerns about taking Entocort, ask your doctor or pharmacist.

Keep this leaflet with your Entocort.

You may need to read it again.

What ENTOCORT is for

Entocort is used in the treatment of Crohn's disease. It can be used to treat acute attacks.

Crohn's disease is an inflammatory disease of the bowel. It mainly affects the small bowel and the first part of the large bowel and causes symptoms such as stomach pain, diarrhoea and fever.

Entocort contains budesonide. This belongs to the group of medicines called corticosteroids, which are used to help reduce inflammation in many parts of the body.

Entocort capsules are designed to release their contents in the small

bowel and the first part of the large bowel.

Ask your doctor if you have any questions about why Entocort has been prescribed for you.

Your doctor may have prescribed it for another reason.

Entocort is not addictive.

Entocort is available only with a doctor's prescription.

Before you take ENTOCORT

When you must not take it

Do not take Entocort if you have an allergy to:

- any medicine containing budesonide
- any ingredients listed at the end of this leaflet.

Some of the symptoms of an allergic reaction may include:

- rash, itching or hives on the skin
- shortness of breath, wheezing or difficulty breathing
- swelling of the face, lips, tongue or other parts of the body.

Do not give Entocort to children.

There is no information available on its use in children.

Do not take Entocort after the expiry date (EXP) printed on the pack or if the packaging is torn or shows signs of tampering.

If it has expired or is damaged, return to your pharmacist for disposal.

If you are not sure whether you should start taking Entocort, talk to your doctor or pharmacist.

Before you start to take it

Tell your doctor if you have allergies to any other medicines, foods, preservatives or dyes.

Tell your doctor if you have or have had any medical conditions, especially the following:

- any recent infection (including chicken pox and measles)
- tuberculosis
- diabetes
- liver problems
- stomach ulcers
- osteoporosis
- hypertension (high blood pressure)
- eye problems (such as glaucoma or cataracts)

It may not be safe for you to take Entocort if you have any of these conditions.

Tell your doctor if you have NOT had chicken pox or measles.

These diseases may be more serious if you get them while taking Entocort. Your doctor may want to vaccinate you for them before you start on Entocort.

Do not take Entocort if you are pregnant or breastfeeding unless your doctor says so. Ask

your doctor about the risks and benefits involved.

It is not known if it is safe for you to take Entocort while you are pregnant. It may affect your baby.

Like most corticosteroid medicines, it is recommended that you do not breastfeed while taking Entocort as it is found in breast milk.

If you have not told your doctor about any of the above, tell them before you start taking Entocort.

Taking other medicines

Tell your doctor or pharmacist if you are taking any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and Entocort may interfere with each other. These include:

- other corticosteroid medicines such as tablets, asthma inhalers, nasal sprays, eye/nose drops
- medicines used to treat fungal infections (eg ketoconazole)
- cimetidine, a medicine used to treat reflux and stomach ulcers

These medicines may be affected by Entocort or may affect how well it works. You may need different amounts of your medicines, or you may need to use different medicines. Your doctor or pharmacist will advise you.

Your doctor and pharmacist have more information on medicines to be careful with or avoid while taking Entocort.

How to take ENTOCORT

Follow all directions given to you by your doctor carefully.

They may differ from the information contained in this leaflet.

How much to take

The usual dose is 3 capsules taken once daily in the morning, before breakfast.

How to take it

Swallow Entocort capsules whole with a glass of water. Do not crush or chew the capsules. If the granules are chewed or crushed they will not work properly.

How long to take it

Continue taking Entocort for as long as your doctor tells you. A treatment course should normally not exceed 12 weeks.

Entocort should not be stopped suddenly. The dose should be reduced gradually over the last 2 to 4 weeks.

If you forget to take it

If you miss a dose, take it as soon as you remember, and then go back to taking it as you would normally.

If it is almost time for your next dose, skip the dose you missed and take your next dose when you are meant to.

Do not take a double dose to make up for the dose that you missed.

This may increase the chance of you getting an unwanted side effect.

If you are not sure what to do, ask your doctor or pharmacist.

If you have trouble remembering when to take your medicine, ask your pharmacist for some hints.

If you take too much (overdose)

Immediately telephone your doctor, pharmacist, the Poisons Information Centre (131 126) or go to Accident and Emergency at your nearest hospital if you think that you or anyone else may have taken too much Entocort. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are taking ENTOCORT

Things you must do

Take Entocort exactly as your doctor has prescribed.

If you have not had chicken pox or measles, avoid close contact with anyone who has these diseases while you are taking Entocort.

Tell your doctor straight away if you think that you have been exposed to chicken pox or measles.

Tell your doctor if you have an infection while you are taking Entocort.

It may not be safe for you to continue taking Entocort if you have an infection.

If you are about to be started on any new medicine, remind your doctor or pharmacist that you are taking Entocort.

Tell any other doctors, dentists and pharmacists who are treating you that you are taking Entocort.

If you become pregnant while using Entocort, tell your doctor or pharmacist immediately.

Things you must not do

Do not eat grapefruit or drink grapefruit juice while you are taking Entocort.

Grapefruit juice, but not other fruit juices, can affect Entocort levels in

the body. This may increase the chance of getting unwanted side effects.

Do not stop taking Entocort unless your doctor tells you to.

If you need to stop taking Entocort, your doctor will tell you how to do it gradually.

Do not use it to treat any other complaints unless your doctor tells you to.

Do not give this medicine to anyone else.

Side effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking Entocort.

All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects.

Ask your doctor or pharmacist to answer any questions you may have.

Tell your doctor or pharmacist if you notice any of the following and they worry you:

- indigestion, flatulence
- nausea, vomiting, abdominal pain
- rapid heart rate or palpitations
- headache, dizziness
- swelling/rounding of the face, weight gain
- back pain
- muscle cramps
- tiredness
- trouble sleeping
- tremor, feeling nervous
- mood swings
- menstrual problems
- skin rash

These side effects are usually mild.

If your medicine has been changed from other oral corticosteroids tablets (eg. prednisone, prednisolone or methylprednisolone) to Entocort you may notice some symptoms that bothered you earlier, eg rash, or pain in muscles and joints. If this happens or you get headaches, nausea or vomiting, or feel tired please contact your doctor.

Tell your doctor or pharmacist if you notice any of the following:

- sign or symptoms of an infection
- blurred vision

These may be serious side effects. You may need medical attention.

Tell your doctor immediately or go to Accident and Emergency at your nearest hospital if you notice any of the following:

- shortness of breath, wheezing or difficulty breathing
- swelling of the face, lips, tongue or other parts of the body
- severe rash

These are very serious side effects. You may need urgent medical attention or hospitalisation. These side effects are very rare.

Tell your doctor or pharmacist if you notice anything else that is making you feel unwell.

Other side effects not listed above may occur in some patients.

If you have changed from or used high doses of oral corticosteroids over a long period of time, your adrenal glands may be affected. Your doctor may do tests to check how the adrenal glands are working.

Your doctor may also tell you to take additional oral corticosteroids during periods of stress such as trauma and surgery.

Ask your doctor to answer any questions you may have.

After taking ENTOCORT

Storage

Keep Entocort in a cool dry place where the temperature stays below 30°C.

Keep your Entocort capsules in the bottle until it is time to take them. Replace the cap firmly after use.

If you take Entocort out of the bottle it will not keep well.

Do not store Entocort or any other medicine in the bathroom or near a sink. Do not leave it on window sills or in the car.

Heat and dampness can destroy some medicines.

Keep it where young children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Disposal

If your doctor tells you to stop taking Entocort or it has passed its expiry date, ask your pharmacist what to do with any capsules you have left over.

Product description

What it looks like

Entocort capsules are two-piece hard gelatin capsules, with an opaque light grey body and an opaque pink cap. The capsule is printed with 'CIR 3 mg' in black.

Entocort is supplied in bottles containing 90 capsules.

Ingredients

Each Entocort capsule contains 3 mg of budesonide as the active

ingredient, and the following
inactive ingredients:

- ethylcellulose
- tributyl acetylcitrate
- methacrylic acid copolymer
- triethyl citrate
- dimethicone
- Polysorbate 80
- purified talc [E553(b)]
- sugar spheres

Each capsule is made from gelatin (E441), with colouring agents titanium dioxide (E171) and iron oxide (E172).

Supplier

AstraZeneca Pty Limited
ABN 54 009 682 311
Alma Road
North Ryde NSW 2113

This leaflet was prepared in March 2009.

Australian Registration Number:

Entocort capsules: AUST R 62812

®Entocort is a trademark of the AstraZeneca group of companies.